



Integrating experimental (in vitro and in vivo) neurotoxicity studies of low-dose thimerosal relevant to vaccines.

Dórea JG¹.

Author information

Abstract

There is a need to interpret neurotoxic studies to help deal with uncertainties surrounding pregnant mothers, newborns and young children who must receive repeated doses of Thimerosal-containing vaccines (TCVs). This review integrates information derived from emerging experimental studies (in vitro and in vivo) of low-dose Thimerosal (sodium ethyl mercury thiosalicylate). Major databases (PubMed and Web-of-science) were searched for in vitro and in vivo experimental studies that addressed the effects of low-dose Thimerosal (or ethylmercury) on neural tissues and animal behaviour. Information extracted from studies indicates that: (a) **activity of low doses of Thimerosal against isolated human and animal brain cells was found in all studies and is consistent with Hg neurotoxicity**; (b) **the neurotoxic effect of ethylmercury has not been studied with co-occurring adjuvant-AI in TCVs**; (c) **animal studies have shown that exposure to Thimerosal-Hg can lead to accumulation of inorganic Hg in brain**, and that (d) **doses relevant to TCV exposure possess the potential to affect human neuro-development**. Thimerosal at concentrations relevant for infants' exposure (in vaccines) is toxic to cultured human-brain cells and to laboratory animals. The persisting use of TCV (in developing countries) is counterintuitive to global efforts to lower Hg exposure and to ban Hg in medical products; its continued use in TCV requires evaluation of a sufficiently nontoxic level of ethylmercury compatible with repeated exposure (co-occurring with adjuvant-AI) during early life.

PMID: 21350943 DOI: [10.1007/s11064-011-0427-0](https://doi.org/10.1007/s11064-011-0427-0)

[Indexed for MEDLINE]

Publication types, MeSH terms, Substances

LinkOut - more resources